

Emerging spectra of singular correlation matrices under small power-map deformations

Vinayak,^{1,*} Rudi Schäfer,^{2,†} and Thomas H. Seligman^{1,3,‡}

¹*Instituto de Ciencias Físicas, Universidad Nacional Autónoma de México, C.P. 62210 Cuernavaca, México*

²*Faculty of Physics, University of Duisburg-Essen, Germany*

³*Centro Internacional de Ciencias, C.P. 62210 Cuernavaca, México*

(Dated: April 19, 2013)

A spectral study of power-map deformed singular random correlation matrices is presented in this paper. The study is inspired from the multivariate analysis of large dimensional datasets where stationarity or quasi-stationarity could often be a good approximation for a non-stationary time series only over a short time horizon. Correlation matrices estimated over such short time horizons are singular, with a significant portion of the spectrum falling exactly on zero. The power map has been proposed to remove this spectral degeneracy, where the correlation matrix is deformed as soon as the exponent is varied from one. Importance of the power map has been illustrated for the singular matrices encountered in financial application, viz. the portfolio optimization. To get a clearer picture of deformed matrices, the exponent of the power map has been chosen to be very close to one such that the emerging spectrum corresponding to the zeros of the original spectrum falls into a gap between zero and nonzero part of the entire spectrum. Analytical approximate estimate of the first and the second moments for the emerging spectra have been calculated for the deformed Wishart matrices of random matrix theory. Furthermore, the study has been extended to the correlated Wishart ensembles with two examples, viz. when the true correlation matrix has block-diagonal structure while in the second example it has a smooth band profile off the diagonal. In both cases the spectra have been analyzed numerically using the analytical results as a benchmark situation. It is shown that the emerging spectra are sensitive to the true correlation.

PACS numbers: 02.50.Sk, 05.45.Tp, 89.90.Gh, 89.65.+n

I. INTRODUCTION

Physics has long concentrated on equilibrium states, but more recently stationary states have drawn increasing attention, particularly in all kinds of transport ranging from heat over charge and particle transport all the way to traffic models [1–4]. Yet it has become very clear that stationarity is at best an approximation for fairly short times in many applications. In this context analysis of financial markets have been in the forefront [5–9], and today the new frontier is the analysis of non-stationary systems with a hope that it can be extrapolated from quasi-stationarity over shorter time periods. A nice example thereof is a recent paper on market states [10].

Starting with early work in biology, multivariate analysis [11, 12] has been established as an essential tool. In particular, correlation matrices play an important role in many problems of biology [13, 14], medical research [15], econophysics [16–18], etc. The eigenlevel-eigenvector analysis of such positive definite matrices has been remarkably useful in the time series analysis of multidimensional complex systems [19–26]. Usually in such studies the time horizon is considered to be longer than the dimensionality of datasets to get rid of spectral redundancies.

Random matrices were introduced by Wishart [27] as a fundamental tool in multivariate analysis. After their successful applications in nuclear and atomic physics [28, 29] random matrices achieved considerable attention from physicists which resulted in the foundation of random matrix theory (RMT). Consequently, RMT has been extensively used and developed in diverse areas of research, for example, in physics [30, 31], in communication engineering [32] and in econophysics [33]. Wishart introduced random matrices of type $C = AA^t/T$ where the matrix A is $N \times T$, A^t is the transpose of A and matrix elements A_{jk} are independent Gaussian variables with mean zero and variance one. In this model C characterizes a *null hypothesis* where the covariance matrix is calculated, over stochastic time series of time horizon T , for N statistically independent variables. In random matrix theory (RMT) ensembles of such matrices are well studied and known as Wishart or Laguerre ensembles (WE) [28]. Analogous to WE the correlated Wishart ensembles (CWE) are defined where matrix elements of A can have genuine nonrandom 'true correlations' [11]. Recently, important advances have been made for CWE [34–40] which go beyond the null hypothesis and set a theoretical background for incorporating the true correlations. Ongoing research activities on time series analysis of complex systems often exploit results of such random matrix models, mainly because these models supply a framework to filter signals from noise or for quantifying the noise [19–24, 26]. While the eigenlevel statistics, e.g. density of eigenlevels, have been useful for denoising

*Electronic address: vinayaksps2003@gmail.com

†Electronic address: rudi.schaefer@uni-due.de

‡Electronic address: seligman@ce.fis.unam.mx

the spectrum, the eigenvector analysis has been useful in identifying the collective movements in the datasets [19, 20].

For short time series, such as we necessarily obtain for non-stationary or quasi-stationary situations, often the time horizon is smaller than the dimensionality of datasets to be analyzed. In that case, because of the factorisable structure, $\mathbf{C}$ will be a singular correlation matrix with at least $N - T$ zero eigenlevels. The corresponding random matrix ensemble is sometimes called "Anti-Wishart ensemble" [41, 42]. For such matrices an analysis of the eigenvectors can only be fruitful for the nonzero eigenlevels, as those of the zero eigenlevels are only defined up to an orthogonal transformation in the degenerate subspace. However, we propose to remove such degeneracy by using the so-called power map which has been successful for regular, or full rank, correlation matrices encountered in financial applications [43–45]. Interestingly, this method produces significantly useful results for singular $\mathbf{C}$, as well, as we show in this paper. In the present paper we are not so much interested in the noise reduction aspect of the power map, but in the breaking of the degeneracy. We will get a much clearer picture by using the exponent values sufficiently close to one so that the part of the spectrum originating from the originally-zero eigenlevels will fall entirely into the gap between zero and the bulk of the spectrum which is close to the spectrum of the correlation matrix. We thus obtain a disjoint spectrum of eigenlevels which we shall call the *emerging spectrum*.

We derive analytically approximate estimate for the first two moments, for the density of emerging eigenlevels and also for the density of the corrections to non-zero eigenlevels of $\mathbf{C}$, in a linear response regime. In analytical work we mostly consider WE because WE is not only analytically more tractable than CWE but it also belongs to a benchmark situation or the null hypothesis, as mentioned above. Besides, we study spectral statistics for power map deformed correlation matrices corresponding to CWE. We show that the true cross-correlations change the spectral statistics appreciably. For CWE we consider two types of ensemble or true correlation matrices. The first one correspond to the 'factor models' which have been useful in quantitative finance [46] and the other one has a smooth band profile.

The paper is organized as follows. In Sec. II we illustrate the power map for singular correlation matrices and describe its importance for the portfolio optimization. In Sec. III we summarize useful results known for WE and for CWE, specially those which we use in this paper. In Sec. IV, we derive estimates of the first and the second moment of the density of small eigenlevels of the power mapped Wishart matrices. In Sec. V we consider non-vanishing true cross correlation in $\mathbf{C}$, for CWE, and analyze the spectra of power mapped matrices. This is followed by conclusion in Sec. VI.




FIG. 1: Portfolio variance Ω^2 normalized to the minimal portfolio variance Ω_0^2 . The length T of the time series was varied, the number of companies was fixed at $N = 100$. Results are shown for sample correlations without noise reduction (black dashed-dotted line) and for power-mapped correlations (blue solid line). The red dashed line at $\Omega^2 / \Omega_0^2 = 4.37$ corresponds to a homogeneous portfolio with all weights equal to $1/N$.

II. POWER MAP FOR SINGULAR CORRELATION MATRICES

We wish to briefly illustrate the power map for singular correlation matrices. For instance, following Markowitz [16], we consider a portfolio of N stocks and wish to calculate the portfolio weights w_{opt} which minimize the portfolio variance

$$\Omega^2 = w_{\text{opt}}^t \Sigma w_{\text{opt}}. \quad (1)$$

In a model setting, we can calculate the minimal variance portfolio Ω_0^2 using the model covariance matrix Σ_0 . In practice, however, the covariance matrix has to be estimated using historical data of finite length T . The shorter the length T of the time series, the noisier is the covariance estimation. Using noisy covariance matrices for portfolio optimization leads to very bad results, see Fig. 1. In this case, the portfolio variance Ω^2 increases as $(1 - N/T)^{-1}$, in accordance with literature [47]. Clearly, it is necessary to improve the estimation of the covariance matrix to obtain better results. The variances of the single stocks can be estimated rather well on short time horizons due to a slowly decaying autocorrelation. The noise in the correlation coefficients C_{kl} can be reduced effectively using the power map

$$C_{kl}^{(q)} = \text{sign } C_{kl} |C_{kl}|^q. \quad (2)$$

The results are presented in Fig. 1. The power map yields portfolio variances which are well below the homogeneous portfolio, even for $N > T$ where the correlation matrix becomes singular. The values for q used in this study range from 1.1 to 2.4. For details of the simulation we refer to Appendix A.




FIG. 2: Density of eigenlevels of $C^{(q)}$ where $q = 1.001$, $N = 1024$ and $\kappa = 1/2$. In Fig. 2(a) we show the density of emerging spectra while in Fig. 2(b) we show the density of the non-zero eigenlevels which is closely described by the Marcenko Pastur law (6). Both densities are shown on different scales. Density in Fig. 2(a) is normalized to $1 - \kappa$ while density in Fig. 2(b) is normalized to κ . Note that the density in Fig. 2(a) is not symmetric, as it gives an impression of some kind of semi-circle. The problem dealt in this paper addresses to the moments of the density in Fig. 2(a).

As far as spectral statistics is concerned, we note that $C^{(q)}$ is always Hermitian, but for $q \neq 1$ it is not necessarily a positive definite matrix in general. For short time horizons, when q is close to 1, the density of the eigenlevels of $C^{(q)}$ is described on two cuts, one which is close to 0 and the other is close to the cut which describes the density of the eigenlevels of C . The former comes because of the spontaneous breaking in the degeneracies of the zero eigenlevels of C . For example, for WE when $q = 1.001$, we show in Fig. 2(a) the density of the emerging spectra while in Fig. 2(b) we show the density of the eigenlevels falling near to the non-zero eigenlevels of C . As shown in this figure, the latter is closely described by theory of WE which is known as the Marcenko Pastur law (6) and is represented by solid a line in the figure. These two cuts merge eventually to a single cut, as the power of q is raised. In this paper we address to the statistical properties of the density shown in Fig. 2(a), viz. we derive an estimate of the first two moments of the density.

III. RANDOM CORRELATION MATRICES: WISHART AND CORRELATED WISHART ENSEMBLES

In RMT one defines basically three invariant Wishart ensembles [28]. In this paper we consider only Wishart orthogonal ensemble (WOE). Similarly we consider CWOE among the three invariant CWEs. For WOE the matrix elements of A are real independent Gaussian variable with mean zero and a fixed variance σ^2 . By construction C is a real symmetric semi-positive definite ma-

trix. Using the Gaussian probability measure, the joint probability density (jpd) of the matrix elements of A can be written as

$$P(A) dA \propto \exp \left[-\text{Tr} \frac{AA^t}{2\sigma^2} \right] dA, \quad (3)$$

where dA is infinitesimal volume in $N \times T$ matrix element space. For $T \geq N$, jpd of the matrix elements of C is defined in $N(N+1)/2$ -dimensional matrix element space [11, 27], as

$$P(C) dC \propto (\det C)^{[N(\kappa-1)-1]/2} \exp \left[-\frac{T}{2\sigma^2} \text{Tr} C \right] dC, \quad (4)$$

where $\kappa = T/N$ and $dC = \prod_{j>k}^N dC_{jk} \prod_{j=1}^N dC_{jj}$ is infinitesimal volume in the matrix element space. The jpd of eigenlevels of C , λ_j , for $j = 1, \dots, N$, may be obtained by transforming the variables to eigenlevel-eigenvector space and then integrating over the eigenvectors. It eventually leads to

$$P(\lambda_1, \dots, \lambda_N) \propto \prod_{j=1}^N w(\lambda_j) \prod_{j>k}^N |\lambda_j - \lambda_k|, \quad (5)$$

where $w(\lambda) = \lambda^{[N(\kappa-1)-1]/2} \exp[-N\kappa\lambda/2\sigma^2]$ is the weight function of the associated Laguerre polynomials. We use bar to denote ensemble averages as in the above equation, while for spectral averages we use angular brackets, e.g. $\langle \cdot \rangle = \{\text{tr} \cdot\}/N$. For $\kappa \leq 1$, jpd of the matrix elements and the jpd of the eigenlevels have been derived in [41, 42], where again Laguerre weight functions describe jpd of the latter. Eigenvalue statistics of the WE are known in great detail [28]. The global statistics, i.e. the eigenlevel density, as well as the local statistics, i.e. the n -point correlation functions for $n \geq 2$, of the eigenlevels are known in terms of Laguerre polynomials. However, for $N, T \rightarrow \infty$ such that $\kappa \geq 1$ and finite, the eigenlevel density converges to the Marcenko Pastur law [48]:

$$\bar{\rho}(\lambda) = \kappa \frac{\sqrt{(\lambda_+ - \lambda)(\lambda - \lambda_-)}}{2\pi\sigma^2\lambda}, \quad (6)$$

where $\lambda_{\pm} = \sigma^2(\kappa^{-1/2} \pm 1)^2$ are the end points of the density. Note that for the singular case, i.e. $\kappa < 1$, $\bar{\rho}(\lambda)$ in the above equation is normalized to κ and not to 1. Therefore, taking into account of $(N-T)$ zeros, for $\kappa \leq 1$ we write

$$\bar{\rho}(\lambda) = \kappa \frac{\sqrt{(\lambda_+ - \lambda)(\lambda - \lambda_-)}}{2\pi\sigma^2\lambda} + (1 - \kappa)\delta(\lambda). \quad (7)$$

For large matrices, higher order spectral correlations or the n -point correlation functions, for $n \geq 2$, are the same as those for Gaussian orthogonal ensemble (GOE) [28], therefore WE spectra have universal spectral fluctuations [49].

In the case of true correlations among the matrix elements A_{jk} , one defines CWE [11]. For instance, in order

to take account of the correlation in the rows of $\mathbf{A}$ matrix one defines $\mathbf{C} = \xi^{1/2} \mathbf{B} \mathbf{B}^\dagger \xi^{1/2}$ where the matrix ξ is fixed positive definite matrix which takes account of true correlations and matrix elements of $\mathbf{B}$ are independent Gaussian variables with zero mean and a fixed variance, just as A_{jk} defined for the WE. We remark that in this definition $\overline{\mathbf{C}} = \sigma^2 \xi$ while $\overline{\mathbf{A}^\dagger \mathbf{A}}/N$ is an identity matrix of dimension $T \times T$. Analytical treatment to CWE [34?–37] is much more difficult and involved than to WE. However, some results are known in this case as well. For instance, analogous to the Marčenko Pastur result, resolvent or Stieltjes transform of the density is known as

$$\overline{G}(z) = \left\langle \frac{1}{z - \frac{\sigma^2}{\kappa} (\kappa - 1 + z \overline{G}(z)) \xi} \right\rangle. \quad (8)$$

The density $\overline{\rho}(\lambda)$ can be determined uniquely via inverse transform

$$\overline{\rho}(\lambda) = \mp \frac{1}{\pi} \Im \overline{G}(\lambda \pm i\epsilon), \quad (9)$$

for positive infinitesimal ϵ . For finite N and T the result for the density has been recently obtained [?] by using the supersymmetric method. For higher order spectral correlations, only asymptotic result for the two-point function is known [37]. Spectral fluctuations of CWE are not always universal rather there are deviations in some cases, as noted for the number variance [29], $\Sigma^2(r)$, for large spectral correlation length r . Besides, for some models of the true correlation matrix ξ few eigenlevels are found to be separated from the bulk density. These eigenlevels often show collective behaviour, therefore, referred to as collective modes. A detail discussion of the spectral properties of CWE can be found in [37].

IV. ESTIMATION OF THE MOMENTS OF EMERGING SPECTRA IN LINEAR RESPONSE REGIME

We consider the power map for the correlation matrices which correspond to the WOE. As mentioned above, we use a power map in order to break the factorisable structure of $\mathbf{C}$ while it is sufficient to consider q very close to 1. We may expand the power map, for small $\alpha = (q-1)$, as

$$\begin{aligned} C_{jk}^{(\alpha)} &= C_{jk}^{(0)} \exp \left[\frac{\alpha}{2} \ln[(C_{jk}^{(0)})^2] \right] \\ &= C_{jk} + \frac{\alpha}{2} C_{jk} \ln(C_{jk}^2) [1 + \mathcal{O}(\alpha)], \end{aligned} \quad (10)$$

where in the second equality we drop the superscript, using $\mathbf{C}$ for $\mathbf{C}^{(0)}$ onwards. Next we expand the eigenlevels $\lambda_j(\alpha)$ of $\mathbf{C}^{(\alpha)}$, as

$$\lambda_j(\alpha) = \lambda_j(0) + \alpha (\delta\lambda_j) [1 + \mathcal{O}(\alpha)]. \quad (11)$$

where the $\lambda_j(0)$'s are the eigenlevels of the covariance matrix $\mathbf{C}$, for $j = 1, \dots, N$ and the $\alpha(\delta\lambda_j)$'s are the leading order corrections coming from the power mapping. For short time horizon, i.e. $T < N$, $\lambda_j(0) = 0$ for $j \leq N - T$. For small α , we assume that the statistics of the relative changes in the eigenlevels is dominated by the linear term. Bearing this assumption in mind, analytically, we derive estimates for the moments of the $\alpha(\delta\lambda_j)$'s, i.e. in other word, in the linear response regime. We refer to $\alpha(\delta\lambda_j)$ as the eigenlevels of the emerging spectra, for $j \leq N - T$, otherwise as non-zero eigenlevel corrections. The term emerging spectra should not be confused with the small eigenlevels of $\mathbf{C}$ for $\kappa = 1$. As a matter of fact there are no such emerging spectra for $\kappa \geq 1$. Without loss of generality we consider $\alpha > 0$.

Use of the perturbation theory for the individual eigenlevels is far too complicated in our case. Since we are mainly concerned about spectral statistics, it is sufficient to work in the matrix element space. For integer $n \geq 0$, we define the moments as

$$\begin{aligned} \overline{m}_n &= \frac{1}{N} \overline{\text{Tr} [\mathbf{C}^{(\alpha)}]^n} = \frac{1}{N} \overline{\sum_{j=1}^N (\lambda_j(\alpha))^n}, \\ \overline{m}_{n}^{\text{mp}} &= \frac{1}{N} \overline{\text{Tr} [\mathbf{C}]^n} = \frac{1}{N} \overline{\sum_{j=1}^N (\lambda_j(0))^n}, \end{aligned} \quad (12)$$

where the superscript {mp} denotes the moments which are closely described by the Marčenko Pastur density (6). Next, for the correction eigenlevels we define the moments, as

$$\overline{\delta m}_n \equiv \frac{\alpha^n}{N} \overline{\sum_{j=1}^N (\delta\lambda_j)^n}. \quad (13)$$

In linear response regime $\overline{\delta m}_n$ may also be estimated by

$$\overline{\delta m}_n \simeq \alpha^n \text{ term in } \frac{1}{N} \left[\overline{\text{Tr} (\mathbf{C}^{(\alpha)})^n} - \overline{\text{Tr} (\mathbf{C})^n} \right]. \quad (14)$$

Defining $\overline{\delta m}_n^{(0)}$ as moments of emerging spectra and $\overline{\delta m}_n^{\text{mp}}$ as the moments of non-zero eigenlevel corrections, we write

$$\overline{\delta m}_n = \overline{\delta m}_n^{(0)} + \overline{\delta m}_n^{\text{mp}}. \quad (15)$$

Note that $\overline{\delta m}_n^{\text{mp}} \equiv \overline{\delta m}_n$ for $T \geq N$.

It turns out that in order to calculate the averages in (14) one needs the moments of C_{jk} . Using the jpd of A_{jk} , given in (3), we calculate,

$$\begin{aligned} \overline{(C_{jk})^{2n}} &= \delta_{jk} \left(\frac{2}{T} \right)^{2n} \frac{\Gamma(2n + T/2)}{\Gamma(T/2)} \\ &\quad + (1 - \delta_{jk}) \frac{\Gamma(2n + 1)}{T^{2n} \Gamma(n + 1)} \frac{\Gamma(T/2 + n)}{\Gamma(T/2)} \\ \overline{(C_{jk})^{2n+1}} &= \delta_{jk} \left(\frac{2}{T} \right)^{2n+1} \frac{\Gamma(2n + 1 + T/2)}{\Gamma(T/2)}, \end{aligned} \quad (16)$$

for integer $n \geq 0$. For completeness, we have included details of this calculation in the Appendix B. Using now Eq. (14) and Eq. (16), we calculate the first moment of the correction eigenlevels, as

$$\begin{aligned}\overline{\delta m_1} &= \frac{\alpha}{2N} \sum_{j=1}^N \overline{C_{jj} \log[(C_{jj})^2]} \\ &= \frac{\alpha}{2N} \sum_{j=1}^N \frac{d}{dh} \overline{(C_{jj})^{2h+1}} \Big|_{h=0}.\end{aligned}\quad (17)$$

In the second equality of the above equation we use Eq. (16) for integer $h \geq 1$ and then we take the derivative for continuous h . From Eq. (17) we obtain

$$\overline{\delta m_1} = \alpha \left[\log\left(\frac{2}{T}\right) + \Psi(1 + T/2) \right], \quad (18)$$

where $\Psi(x)$ is the digamma function. Similarly for the second moment we obtain

$$\begin{aligned}\overline{\delta m_2} &= \alpha^2 \left(1 + \frac{2}{T}\right) \left\{ \left(\log\frac{2}{T} + \Psi(2 + T/2) \right)^2 \right. \\ &\quad \left. + \Psi'(2 + T/2) \right\} + \frac{\alpha^2}{\kappa} \left(1 - \frac{1}{N}\right) \left\{ \left(-\log(T) \right. \right. \\ &\quad \left. \left. + 1 - \frac{\gamma}{2} + \frac{\Psi(1 + T/2)}{2} \right)^2 + \frac{\Psi'(1 + T/2)}{4} \right. \\ &\quad \left. \left. - 1 + \frac{\pi^2}{8} \right\},\end{aligned}\quad (19)$$

where $\Psi'(x) = d\Psi(x)/dx$. Up to the leading order, the large T asymptotic of $\overline{(\delta m_1)}$ and $\overline{(\delta m_2)}$ may be evaluated as

$$\begin{aligned}\overline{\delta m_1} &\sim \frac{\alpha}{T}, \\ \overline{\delta m_2} &\sim \frac{\alpha^2}{4\kappa} ([\log(T) + c_1]^2 + c_2),\end{aligned}\quad (20)$$

where $c_1 = \gamma + \log(2) - 2 = -0.729637\dots$ and $c_2 = \pi^2/2 - 4 = 0.934802\dots$, for γ being the Euler constant. Note that the density defined by $\overline{\delta m_n}$ is doubly peaked for $T < N$, one which corresponds to the eigenlevels of emerging spectra and the other corresponds to the non-zero eigenlevel corrections. The two peaks converge to one as soon as $T \geq N$.

As we know from the Marčenko Pastur result that the moments of the density depend on κ alone. While moments defined in (15) may have dependency on T also, as we show below, although to estimate either of the moments is in no way straight forward. However, in the linear response regime we may assume that the ensemble averaged density of the non-zero correction eigenlevels could be approximated by a rescaled Marčenko Pastur density. Since non-negativity of eigenlevels is no longer a valid criterion, we allow the scale to assume negative values as well. We note that this density should be normalized




FIG. 3: Comparison of theory, shown by solid line, with simulations, shown by symbols where T is the variable. In Figs. 3(a) and 3(b) we compare first two moments $\overline{\delta m_1}$ and $\overline{\delta m_2}$ obtained from the numerical simulations with the theory (20). In Figs. 3(c) and 3(d) we compare numerical results of scaling and shifting parameters s and r , obtained from Eqs. (24, 25) with theory (26). In Figs. 3(e) and 3(f) we compare numerical results of $\overline{\delta m_1^{\text{mp}}}$ and $\overline{\delta m_2^{\text{mp}}}$ with theory (27). Finally, in Figs. 3(g) and 3(h) we compare $\overline{\delta m_1^{(0)}}$ and $\overline{\delta m_2^{(0)}}$ with the theory 28.

to 1 for $\kappa \geq 1$ and κ otherwise. Besides, it is suggestive to use a shifting parameter. Finally, we make an ansatz for this density,

$$\bar{\rho}_1(\delta\lambda) = \kappa \frac{\sqrt{(\delta\lambda_+ - \delta\lambda)(\delta\lambda - \delta\lambda_-)}}{2\pi(\delta\lambda - r)s}. \quad (21)$$

Here $\delta\lambda_{\mp} = s(\kappa^{-1/2} \pm 1) + r$ while r and s are the shifting and scaling parameters respectively. We fix these parameters by calculating the first two moments of the density. For the above density (21), we obtain

$$\overline{\delta m_1^{\text{mp}}} = \begin{cases} s + r, & \text{for } \kappa \geq 1, \\ s + \kappa r, & \text{for } \kappa \leq 1. \end{cases} \quad (22)$$

Similarly for the second moment we obtain

$$\overline{\delta m_2^{\text{mp}}} = \begin{cases} \left(1 + \frac{1}{\kappa}\right) s^2 - r^2 + 2r\overline{\delta m_1}, & \text{for } \kappa \geq 1, \\ \left(1 + \frac{1}{\kappa}\right) s^2 - r^2\kappa + 2r\overline{\delta m_1^{\text{mp}}}, & \text{for } \kappa \leq 1. \end{cases} \quad (23)$$

Inverting above relations, we write

$$s = \begin{cases} -\sqrt{\left[\overline{\delta m_2} - \overline{\delta m_1}^2\right] \kappa}, & \text{for } \kappa \geq 1, \\ -\sqrt{\overline{\delta m_2^{\text{mp}}} - \frac{\overline{\delta m_1^{\text{mp}}^2}}{\kappa}}, & \text{for } \kappa \leq 1, \end{cases} \quad (24)$$

and

$$r = \begin{cases} \overline{\delta m_1} - s, & \text{for } \kappa \geq 1, \\ \frac{\overline{\delta m_1^{\text{mp}}} - s}{\kappa}, & \text{for } \kappa \leq 1. \end{cases} \quad (25)$$

We remark that the scaling parameter s should be a negative quantity as the power mapping, for $\alpha > 0$, shrinks the density of the $\lambda_j(\alpha)$'s for $N - T < j \leq N$. For $\kappa \geq 1$ the two parameters are known in terms of $\overline{\delta m_1}$ and $\overline{\delta m_2}$ where both moments behave smoothly at $\kappa = 1$. On the other hand, one expects non-smooth behaviour in $\overline{\delta m_1^{\text{mp}}}$ and $\overline{\delta m_2^{\text{mp}}}$ at $\kappa = 1$, as the moments of the emerging spectra should not behave smoothly at $\kappa = 1$. We now use results (20) and (24, 25) to write asymptotic form of s and r for $\kappa \geq 1$. For large T , we obtain N -independent behaviour for both the quantities, e.g.,

$$\begin{aligned} s &\sim -\frac{\alpha}{2} \sqrt{[\log(T) + c_1]^2 + c_2}, \\ r &\sim \alpha \left(\frac{1}{T} + \frac{\sqrt{[\log(T) + c_1]^2 + c_2}}{2} \right). \end{aligned} \quad (26)$$

Exploiting the N -independence in these results, we may extrapolate them for $\kappa \leq 1$. Then the estimation of $\overline{\delta m_1^{\text{mp}}}$ and $\overline{\delta m_2^{\text{mp}}}$ becomes straight forward, e.g.,

$$\begin{aligned} \overline{\delta m_1^{\text{mp}}} &= \kappa \overline{\delta m_1} + s(1 - \kappa), \\ \overline{\delta m_2^{\text{mp}}} &= \kappa \overline{\delta m_2} - \kappa \overline{\delta m_1}^2 + \frac{(\overline{\delta m_1^{\text{mp}}})^2}{\kappa}. \end{aligned} \quad (27)$$

Using (15), estimation of the moments of the emerging spectra become trivial. For large T and $\kappa \leq 1$, one gets

$$\begin{aligned} \overline{\delta m_1^{(0)}} &= -s(1 - \kappa), \\ \overline{\delta m_2^{(0)}} &= s^2(1 - \kappa). \end{aligned} \quad (28)$$

We mention that for small values of κ , discrepancy in our approach becomes effective and such linear response theory fails to work.

In Fig. 3 we have compared our theoretical estimates with numerical results for fixed $N = 512$ and $N = 1024$. For the numerical results we have use

$\overline{\delta m_n} = \overline{[\sum_{j=1}^N (\Delta \lambda_j)^n] / N}$ where $\Delta \lambda_j = \lambda_j^{(\alpha)} - \lambda_j$. Similarly, for $\overline{\delta m_n^{(0)}}$ and $\overline{\delta m_n^{\text{mp}}}$ we use $\Delta \lambda_j$ for $0 < j \leq N - T$ and $N - T < j \leq N$, respectively. As shown in the graph that our theory gives reasonable account for the numerical results. The non-smooth behaviour in the moments of non-zero or emerging spectra are well described by our theory. As we see in Figs. 3(c) and 3(d) that the scaling and the shifting parameters are independent of N . For the first moment we see some interesting features in Figs 3(e) and 3(g), viz. (i) the two curves are almost symmetric about zero, (ii) while $\overline{\delta m_1^{\text{mp}}}$ reaches to minimum roughly when $T < N\kappa/4$, $\overline{\delta m_1^{(0)}}$ reaches to the maximum value. The second moment for emerging spectra, on the other hand, increases for small T . This feature, however, can not be captured by our theory as κ is small.

V. POWER MAPPING FOR THE CORRELATED WISHART ENSEMBLES

The CWEs are difficult to handle analytically in general and we mainly analyze the spectra numerically. We consider basically two types of ensemble or true correlation matrix ξ . In the first example, ξ has ℓ block diagonal structures, where N_i being the size of the i 'th block such that $\sum_i^\ell N_i = N$ while for the i 'th block $\xi_{jk} = \delta_{jk} + (1 - \delta_{jk}) c_i$. Matrix elements of the connecting off-diagonal blocks are 0. Such examples qualitatively correspond to the 'factor model' in quantitative finance [43, 44]. In the second example, ξ has smooth band structure, e.g., $\xi_{jk} = c^{|j-k|}$ which may correspond to situations where we expect that spatial correlations fall-off exponentially with the spatial distances. Without loss of generality in both cases we consider $c \geq 0$.

A. Ensemble correlation matrix with block structures

The simplest example for which we can readily obtain analytical results is $\ell = 1$. In this example ξ is a dense matrix and its eigenlevels ξ_j are simply given by $\xi_j = 1 - c$ for $j = 1, \dots, N-1$ and $\xi_N = Nc + 1 - c$. For this spectrum, Eqs. (8, 9) yields the density [37]

$$\begin{aligned} \overline{\rho}(\lambda) &= \overline{\rho'}(\lambda) + \delta \left(\lambda - \frac{(Nc + 1 - c)(Nc\kappa + 1 - c)}{Nc\kappa} \right), \\ \overline{\rho'}(\lambda) &= \kappa \frac{\sqrt{(\lambda_+ - \lambda)(\lambda - \lambda_-)}}{2\pi(1 - c)\lambda}, \end{aligned} \quad (29)$$

where $\lambda_{\pm} = (1 - c)(\kappa^{-1/2} \pm 1)^2$. Note the similarity of $\overline{\rho'}(\lambda)$ with the Marčenko-Pastur law (6), where only a factor $1 - c$ appears extra in the place of σ^2 . The delta function appears in the above result as long as $c \geq (N\sqrt{\kappa})^{-1}$. It is known from the work of Baik *et al* that the density of the separated eigenlevels or the collective mode




FIG. 4: Results for power mapping of correlation matrices corresponding to CWE. We choose the true correlation matrix model where the matrix elements are $\xi_{jk} = \delta_{jk} + c(1 - \delta_{jk})$ and $c = 0.5$. Bulk densities of $\bar{\rho}(\lambda)$, $\bar{\rho}_1(\delta\lambda^{(1)})$ and $\bar{\rho}_0(\delta\lambda^{(0)})$ are shown with their tail in the insets, respectively in Figs. 4(a), 4(b) and 4(c). Theory for the distribution of the isolated eigenlevels which is a Gaussian distribution as shown in Fig. 4(a), is obtained by using results for the mean and the variance given in [37]. We also remark about the existence of collective mode in Fig. 4(c). Distribution of isolated eigenlevels in insets of Figs. 4(b) and 4(c) both are not Gaussian. Averaged mean positions as a function of the correlation coefficient c , for $\kappa = 1/2$, are shown in Figs. 4(d), 4(e) and 4(f), respectively for the separated eigenlevels in $\bar{\rho}(\lambda)$, $\bar{\rho}_1(\delta\lambda^{(1)})$ and $\bar{\rho}_0(\delta\lambda^{(0)})$.

is described by a Gaussian distribution [50]. Analytical results for the ensemble averaged mean and the variance are known [37] in terms of the spectra of ξ .

To obtain some of the analytical results for the power mapped correlation matrices of this model we simply use the fact that for the bulk only the variance is reduced and derive results for the moments of bulk densities only. Using $1 - c$ as the variance in Eqs. (B3, B5, B6) and following the calculations of Sec. III, we readily find

$$\begin{aligned} \overline{\delta m_1} &\sim \alpha(1 - c) \ln(1 - c), \\ \overline{\delta m_2} &\sim \overline{\delta m_1}^2 + \frac{\alpha^2(1 - c)^2}{4\kappa} \left([\ln(T) + c_1 - 2\ln(1 - c)]^2 + c_2 \right). \end{aligned} \quad (30)$$

Next, extending the ansatz (21) for the density $\bar{\rho}(\lambda)$, we calculate

$$\overline{\delta m_1^{(1)}} = \begin{cases} s\overline{\delta m_1} + r, & \text{for } \kappa \geq 1, \\ s\overline{\delta m_1} + \kappa r, & \text{for } \kappa \leq 1. \end{cases} \quad (31)$$




FIG. 5: Comparison of theory with numerical results where for the matrix elements of the true correlation matrix are $\xi_{jk} = \delta_{jk} + c(1 - \delta_{jk})$. We choose $c = 0.5$. First and second moments, $\overline{\delta m_1}$ and $\overline{\delta m_2}$ are shown respectively in Figs. 5(a) and 5(b). Scaling and shifting parameters s and r are shown respectively in Fig. 5(c) and Fig. 5(d). Moments of the density of non-zero eigenlevel corrections, viz. $\overline{\delta m_1^{(1)}}$ and $\overline{\delta m_2^{(1)}}$, are shown respectively in Figs. 5(e) and 5(f). Moments of the emerging spectra, viz. $\overline{\delta m_1^{(0)}}$ and $\overline{\delta m_2^{(0)}}$ are shown respectively in Figs. 5(g) and 5(h). Solid lines are theories and open circles represent numerical results. These moments are calculated only for the bulk densities. Dashed lines are theoretical results of the WE.

$$\overline{\delta m_2^{(1)}} = \begin{cases} \overline{m_2}s^2 + r^2 + 2sr\overline{m_1}, & \text{for } \kappa \geq 1, \\ \overline{m_2}s^2 + r^2\kappa + 2sr\overline{m_1} & \text{for } \kappa \leq 1, \end{cases} \quad (32)$$

where $\overline{m_1}$ and $\overline{m_2}$ are the first and the second moments $\bar{\rho}(\lambda)$ which in this case are $1 - c$ and $(1 - c)^2(1 + 1/\kappa)$, respectively and we have replaced the superscript {mp} by (1). For $\kappa \geq 1$ and for a given c , s and t can be quickly estimated in terms of $\overline{\delta m_1}$ and $\overline{\delta m_2}$:

$$\begin{aligned} s &= -\sqrt{\frac{\kappa[\overline{\delta m_2} - \overline{\delta m_1}^2]}{(1 - c)^2}}, \\ r &= \overline{\delta m_1} - s(1 - c). \end{aligned} \quad (33)$$

Since this result confirms possibility of extrapolating s




FIG. 6: Non-zero eigenlevel density and emerging spectra eigenlevel density, of $C^{(q)}$ where the true correlation matrix ξ has ℓ -diagonal blocks and $\ell = 3$, respectively in Figs. 6(a) and 6(b). We have considered $N_1 = N/2$, $N_2 = N_3 = N/4$ while correlation coefficients are respectively $c_1 = 0.9$, $c_2 = 0.45$ and $c_3 = 0.225$. In the inset of Fig. 6(a) density of the isolate eigenlevels are shown with the theory. First and second moments, $\overline{\delta m_1^{(0)}}$, $\overline{\delta m_2^{(0)}}$, of $\bar{\rho}_0(\lambda^{(0)})$ are shown as function of time horizon T , respectively in Fig. 6(c) and in Fig. 6(d). In Fig. 6(a) solid lines represent theory obtained from the numerical solution of Eq. (9) and symbols represent numerical data. For $c = 0$, which corresponds to WE, only theories are shown by dashed line.

and r for $\kappa \leq 1$, therefore for the moment $\overline{\delta m_1^{(1)}}$ we get

$$\overline{\delta m_1^{(1)}} = \kappa \overline{\delta m_1} + (1 - c)s(1 - \kappa), \quad (34)$$

while the result for second moment is the same as obtained in Eq. (27).

In Fig. 4(a), Fig. 4(b) and Fig. 4(c), we show bulk densities for the non-zero eigenlevels, non-zero correction eigenlevels and emerging spectra of $C^{(q)}$, respectively, for $c = 0.5$. Theory for Fig. 4(a) comes from Eq. (29) while in Fig. 4(b) it comes from the ansatz for $\bar{\rho}'_1(\delta\lambda)$ with s and r extrapolated for $\kappa \leq 1$. Theory of Fig. 4(c) is not known. Densities observed in these figures are different from those for the WE. In the inset of these figures we show the density of isolated eigenlevels which is remarkable in Fig. 4(c). In Fig. 4(d), Fig. 4(e) and Fig. 4(f), we show averaged mean of the largest eigenlevel respectively for $\bar{\rho}(\lambda)$, $\bar{\rho}_1(\delta\lambda^{(1)})$ and $\bar{\rho}_0(\delta\lambda^{(0)})$. If the separation is large then the corrections due to the power map to the averaged mean of the largest eigenlevel may be estimated by $\overline{\delta\lambda_N} \sim \alpha \overline{\lambda_N} \log(\lambda_N^2)/2$. In Fig. 4(e) we have compared this theory with numerics. Note that the collective mode that appeared in Fig. 4(b) is in the other side of the peak of the density in bulk, as opposed




FIG. 7: Eigenvalue density, $\bar{\rho}(\lambda)$, of C , emerging spectral density, $\bar{\rho}_0(\lambda^{(0)})$, of $C^{(q)}$ for different values of the correlation coefficient c are shown in Fig. 7(a) and in Fig. 7(b), respectively. First and second moments, $\overline{\delta m_1^{(0)}}$, $\overline{\delta m_2^{(0)}}$, of $\bar{\rho}_0(\lambda^{(0)})$ are shown as function of time horizon T , respectively in Fig. 7(c) and in Fig. 7(d). In Fig. 7(a) solid lines represent theory obtained from the numerical solution of Eq. (9) and symbols represent numerical data. For $c = 0$, which corresponds to WE, only theories are shown by dashed line.

to what is seen in Fig. 4(a) and in Fig. 4(c).

In Fig. 5 we compare the numerical moments with theoretical results. We also show in this figure theoretical results of the WE, those we derived in Sec. IV. The moments of $\bar{\rho}_{(0)}(\delta\lambda^{(0)})$ are observed to be smaller than their WE counterparts. Qualitatively similar results are obtained for $\ell = 3$ where we have considered $N_1 = N/2$, $N_2 = N_3 = N/4$ with correlation coefficients respectively as $c_1 = 0.9$, $c_2 = 0.45$ and $c_3 = 0.225$. Here in Fig. 6(b) we notice a hump in the tail, indicating that the delta functions are about to merge with the bulk.

B. Ensemble correlation matrix with smooth band structure

In our second example, i.e. $\xi_{jk} = c^{|j-k|}$, we present numerical results only. The density of eigenlevels of the correlation matrices is obtained by solving the Eq. (9) numerically for the spectrum of ξ while $\kappa = 1/2$. For different values of correlation coefficient, viz. $c = 0.8, 0.5$ and $c = 0$, densities are compared in Fig. 7 (a) where we find a good agreement of the theory with numerics. As shown in this figure, majority of eigenlevels move towards zero with c while large eigenlevels moves away from the upper edge of Marčenko-Pastur density. In Fig. 7 (b) we compare densities of emerging spectra which change

with c as well. We remark that in this figure the upper and the lower both spectral edges moves efficiently with c because there is no hard edge as such there is in the density of emerging spectra. In Figs. 7 (c) and 7 (d) the first two moments of $\bar{\rho}_0(\lambda^{(0)})$ are compared for different time horizon. These moments qualitatively explain what we see in Fig. 7 (b). Interestingly for strong correlations these moments develop convex profile which worths further investigations.

VI. CONCLUSION

In conclusion, we have studied spectral behaviour of random correlation matrices estimated for short time horizons and deformed slightly by the power map. This study is relevant in the multivariate analysis of large complex systems. In the context of financial time series we have shown the importance of the power map with an illustration of this method for singular correlation matrices. We have mainly concentrated on the statistics of the spectra emerging due to the exponent of the power map being very close to one. On one hand this is interesting due to the spontaneous breaking of the degeneracy at zero while on the other hand we get a clearer picture for the spectral statistics and analytical treatment becomes easier. Our analytic results for the Wishart ensembles set a benchmark which can be important in detecting the true correlations. In the derivation of these results we have introduced scaling and shifting parameters which are calculated for $\kappa \geq 1$. Then by extrapolating these parameters for $\kappa \leq 1$ we have calculated estimates of the first and the second moments of the emerging spectra. These results are valid in a linear response regime and therefore are not valid for very small κ , at least for the higher order moments. We should emphasized that smaller κ values are also important for the real data. Moreover, for a very singular matrix statistical analysis for the emerging spectra suits better than the regular nonzero spectra which has only few eigenlevels. It is therefore important to obtain results for the spectral statistics for very small κ values.

We have also considered true correlations in the matrix elements. We have focused on mainly two types examples, viz. when the ensemble correlation matrix has a block diagonal structure and when it has smooth band structure. For the simplest case of the former model we have been able to obtain theoretical results for the bulk. In that case appearance of collective mode in the density of the emerging spectra is interesting, although analytically it can not be described by using linear response theory. In the second case we have numerically shown that how true correlations change the moments appreciably. With both examples we have established the that the emerging spectra is also very sensitive to correlations. As far as analytic treatment in concerned we believe that some scaling and shifting parameter may be useful in explaining these numerical results. Finally we mention that

it would be worth investigating properties of eigenvectors corresponding to the emerging spectra, as well.

VII. ACKNOWLEDGMENT

We are thankful to F. Leyvraz and T. Guhr for important and useful discussions. We acknowledge financial support from the project 44020 by CONACyT, Mexico, and project PAPIIT UNAM RR 113311, Mexico. Vinayak is a postdoctoral fellow in DGAPA/UNAM.

Appendix A: Details of the portfolio optimization study

The weight vector of the minimal variance portfolio is calculated as

$$w_{\text{opt}} = \frac{\Sigma^{-1}e}{e^t \Sigma^{-1}e}, \quad (\text{A1})$$

where e is a vector of length N with all entries set to one, e^t denotes the transposed vector. In order to calculate the optimal weights (A1), we need to know the covariance matrix Σ of the N stock returns. In practice, this covariance matrix has to be estimated using historical data. Here we consider a model setting with $N = 100$ stocks, a model correlation matrix C_0 with 5 blocks of size 20, corresponding to industry sectors, and randomly distributed but fixed standard deviations σ_k . In a factor model that reflects our correlation matrix C_0 , we simulate time series of length N , estimate the correlation matrix $C^{(\text{samp})}$ and apply the power map to arrive at the matrix $C^{(q)}$ with entries

$$C_{kl}^{(q)} = \text{sign } C_{kl}^{(\text{samp})} |C_{kl}^{(\text{samp})}|^q. \quad (\text{A2})$$

We multiply with the standard deviations to get the elements of the covariance matrix,

$$\hat{\Sigma}_{kl} = \sigma_k \sigma_l C_{kl}^{(q)}. \quad (\text{A3})$$

With this covariance matrix we calculate the weights

$$\hat{w}_{\text{opt}} = \frac{\hat{\Sigma}^{-1}e}{e^t \hat{\Sigma}^{-1}e}. \quad (\text{A4})$$

Using the model correlation matrix C_0 and the corresponding covariance matrix $\Sigma_0 = \sigma C_0 \sigma$, with $\sigma = \text{diag}(\sigma_1, \dots, \sigma_N)$, the actual portfolio variance for the weights (A4) reads

$$\Omega^2 = \hat{w}_{\text{opt}}^t \Sigma_0 \hat{w}_{\text{opt}}, \quad (\text{A5})$$

whereas the minimal variance Ω_0^2 is obtained by calculating the optimal weights (A1) for Σ_0 .

Appendix B: Details of the derivation of Eq.(16)

We begin with considering the following binary associations, for integer $n \geq 1$,

$$\begin{aligned} \overline{(C_{jk})^{2n}} &= \frac{1}{T} \sum_{m_1=1}^T \overline{A_{jm_1} A_{km_1} (C_{jk})^{2n-1}} \\ &+ \frac{1}{T^{2n}} \sum_{i=2}^{2n} \sum_{\{m\}=1}^T \overline{A_{jm_1} A_{km_1} \dots A_{jm_i} A_{km_i} \dots A_{jm_{2n}} A_{km_{2n}}} \\ &+ \frac{1}{T^{2n}} \sum_{i=2}^{2n} \sum_{\{m\}=1}^T \overline{A_{jm_1} A_{km_1} \dots A_{jm_i} A_{km_i} \dots A_{jm_{2n}} A_{km_{2n}}}. \end{aligned} \quad (B1)$$

These binary associations results in terms of the moments of the diagonal elements

$$\begin{aligned} \overline{(C_{jk})^{2n}} &= \delta_{jk} \overline{(C_{jk})^{2n-1}} \left\{ 1 + \frac{2n-1}{T} \right\} + \left(\frac{2n-1}{T} \right) \\ &\times \overline{(C_{jk})^{2n-2} C_{kk}} \\ &= \delta_{jk} \left(1 + \frac{2n-1}{T} \right) \left\{ \sum_{\ell=1}^n \overline{(C_{jk})^{2n-\ell}} \right. \\ &\times \left. \frac{\Gamma(2n+1)\Gamma(n+2-\ell)}{\Gamma(n+1)\Gamma(2n-2\ell+3)(2T)^{\ell-1}} \right\} \\ &+ \frac{\Gamma(2n+1)}{\Gamma(n+1)(2T)^n} \overline{(C_{kk})^n}, \end{aligned} \quad (B2)$$

where we have considered the mean and the variance of A_{jk} to be 0 and 1, respectively. Moments for the diagonal terms are comparatively easier to obtain. Bearing similar binary associations in mind we readily write the result for integer $n > 0$ with arbitrary variance σ^2 of A_{jk} :

$$\begin{aligned} \overline{(C_{jj})^n} &= \prod_{i=1}^n \sigma^2 \left(1 + \frac{2(i-1)}{T} \right) \\ &= \left(\frac{2\sigma^2}{T} \right)^n \frac{\Gamma(n+T/2)}{\Gamma(T/2)}. \end{aligned} \quad (B3)$$

Note here that the second equality in the above equation is the extension of the first equality for $n = 0$. Now we use the result (B3) in Eq. (B2) with $\sigma^2 = 1$ and derive exact result for the even moments of the matrix elements

$$\begin{aligned} \overline{(C_{jk})^{2n}} &= \frac{\Gamma(2n+1)}{\Gamma(n+1)\Gamma(T/2)T^{2n}} \left[\delta_{jk} T \left(1 + \frac{2n-1}{T} \right) \right. \\ &\times \sum_{\ell=1}^n 2^{2(n-\ell)+1} \frac{\Gamma(n+2-\ell)\Gamma(2n-\ell+T/2)}{\Gamma(2n-2\ell+3)} \\ &\left. + \Gamma(n+T/2) \right] \\ &= \frac{\Gamma(2n+1)}{\Gamma(n+1)\Gamma(T/2)T^{2n}} \left[\delta_{jk} T \left(1 + \frac{2n-1}{T} \right) \right. \\ &\times \frac{\Gamma(1/2)\Gamma(2n+T/2) - \Gamma(n+1/2)\Gamma(n+T/2)}{\Gamma(n+1/2)(T+2n-1)} \\ &\left. + \Gamma(n+T/2) \right] \\ &= \frac{\Gamma(2n+1)}{\Gamma(n+1)\Gamma(T/2)T^{2n}} \left[\delta_{jk} \frac{\Gamma(1/2)\Gamma(2n+T/2)}{\Gamma(n+1/2)} \right. \\ &\left. + \Gamma(n+T/2) (1 - \delta_{jk}) \right]. \end{aligned} \quad (B4)$$

On further simplification it results in the first equality of Eq. (16). For arbitrary variance we write Eq. (16) as

$$\begin{aligned} \overline{(C_{jk})^{2n}} &= \delta_{jk} \left(\frac{2\sigma^2}{T} \right)^{2n} \frac{\Gamma(2n+T/2)}{\Gamma(T/2)} + (1 - \delta_{jk}) \\ &\times \left(\frac{\sigma^2}{T} \right)^{2n} \frac{\Gamma(2n+1)}{\Gamma(n+1)} \frac{\Gamma(T/2+n)}{\Gamma(T/2)}. \end{aligned} \quad (B5)$$

Following the similar steps we obtain the second equality of Eq. (16), which is with σ^2 may be written as

$$\overline{(C_{jk})^{2n+1}} = \delta_{jk} \left(\frac{2\sigma^2}{T} \right)^{2n+1} \frac{\Gamma(2n+1+T/2)}{\Gamma(T/2)}. \quad (B6)$$

-
- [1] B. Derrida and J. L. Lebowitz, Phys. Rev. Lett. **80**, 209 (1998); T. H. Seligman and H. A. Weidenmüller, J. Phys. A: Math. Theor. **44**, 205302 (2011); M. S. Green, J. Chem. Phys. **20**, 1281 (1952); R. Kubo, J. Phys. Soc. Japan **12**, 570 (1957).
- [2] C. Maes and K. Netočný, Euro. Phys. Lett. **82**, 30003 (2008).
- [3] K. Nagel and M. Schreckenberg, J. Phys. I France **2**, 2221 (1992); M. Horvat, T. Prosen, G. Benenti and G. Casati, Phys. Rev. E **86**, 052102 (2012).
- [4] B. Buča and T. Prosen, New J. Phys. **14**, 073007 (2012).
- [5] R. N. Mantegna, H. E. Stanley, Physica A **239**, 255 (1997).
- [6] J. Bouchaud, M. Potters, Theory of Financial Risks (Cambridge University Press, Cambridge, 2000).
- [7] J. Voit, The Statistical Mechanics of Financial Markets (Springer, Heidelberg, 2001).
- [8] T. F. Cooley, V. Quadrini, The American Economic Review **91**, 1286 (2001).
- [9] D. Pelletier, Journal of Econometrics **131**, 445 (2006).
- [10] M. C. Münnix, T. Shimada, R. Schäfer, F. Leyvraz, T. H. Seligman, T. Guhr, and H. E. Stanley, Nature Scientific Reports **2**, 644 (2012).
- [11] S. S. Wilks, Mathematical Statistics (Wiley, New York, 1962).
- [12] R. J. Muirhead, Aspects of Multivariate Statistical Theory, Wiley Series in Probability and Statistics (Wiley, 2009).

- [13] I. Lee, D. Yoganarasimha, G. Rao and J. J. Knierim, Nature **430**, 456 (2004).
- [14] J. K. Conner, Nature **420**, 407 (2002).
- [15] H. Johansen-Berg, T. E. J. Behrens, M. D. Robson, I. Drobnjak, M. F. S. Rushworth, J. M. Brady, S. M. Smith, D. J. Higham, and P. M. Matthews, PNAS **101**(36), 13335 (2004).
- [16] H. Markowitz, *Portfolio selection* (John Wiley & Sons, New York, 1959).
- [17] S. Pafka, I. Kondor, Estimated correlation matrices and portfolio optimization, Physica A **343**, 623 (2004).
- [18] A. J. McNeil, R. Frey, P. Embrechts, Quantitative Risk Management: Concepts, Techniques, and Tools, Princeton University Press, 2005, Ch. 4: Financial Time Series.
- [19] L. Laloux, P. Cizeau, J. P. Bouchaud and M. Potters, Phys. Rev. Lett. **83**, 1467 (1999).
- [20] V. Plerou, P. Gopikrishnan, B. Rosenow, L. A. Nunes Amaral and H. E. Stanley, Phys. Rev. Lett. **83**, 1471 (1999).
- [21] V. Plerou, P. Gopikrishnan, B. Rosenow, L. A. Nunes Amaral, T. Guhr, and H. E. Stanley, Phys. Rev. E **65** 066126 (2002).
- [22] F. Lillo and R. N. Mantegna, Phys. Rev. E **72**, 016219 (2005).
- [23] M. S. Santhanam and P. K. Patra, Phys. Rev. E **64**, 016102 (2001).
- [24] F. Luo, J. Zhong, Y. Yang and J. Zhou, Phys. Rev. E **73** 031924 (2006).
- [25] P. Šeba, Phys. Rev. Lett. **91**, 198104 (2003).
- [26] S. Abe and N. Suzuki, arXiv:0909.3830v1 (2009).
- [27] J. Wishart, Biometrika **20A**, 32 (1928).
- [28] M. L. Mehta, *Random Matrices* (Academic Press, New York, 2004).
- [29] T. A. Brody, J. Flores, J. B. French, P. A. Mello, A. Pandey and S. S. M. Wong, Rev. Mod. Phys. **53**, 385 (1981).
- [30] T. Guhr, A. M. Groeling and H. A. Weidenmüller, Phys. Rep. **299**, 189 (1998).
- [31] C. W. J. Beenakker, Rev. Mod. Phys. **69**, 731 (1997).
- [32] R. R. Müller, IEEE Transactions on Information Theory, **48**, 9 (2002); R. R. Müller, Acta Phys. Pol. B **36**, 1001 (2005).
- [33] J. P. Bouchaud and M. Potters, arXiv:0910.1205v1 **q-fin.ST**, (2009).
- [34] A. M. Sengupta and P. P. Mitra, Phys. Rev. E **60**, 3389 (1999).
- [35] S. H. Simon and A. L. Moustakas, Phys. Rev. E **69**, 065101(R) (2004).
- [36] Z. Burda, J. Jurkiewicz and B. Waclaw, Phys. Rev. E **71**, 026111 (2005).
- [37] Vinayak and A. Pandey, Phys. Rev. E **81**, 036202 (2010).
- [38] C. Recher, M. Kieburg, and T. Guhr, Phys. Rev. Lett. **105**, 244101 (2010).
- [39] C. Recher, M. Kieburg, T. Guhr and M. R. Zirnbauer, J. Stat. Phys. **148**, 981 (2012).
- [40] Vinayak and M. Žnidarič, J. Phys. A: Math. Theor. **45**, 125204 (2012).
- [41] S. Maslov and Y.-C. Zhang, Phys. Rev. Lett. **87** 248701 (2001); Y.-K. Yu and Y.-C. Zhang, Physica A: Statistical Mechanics and its Applications **312**, 1 (2002).
- [42] R. A. Janik and M. A. Nowak, J. Phys. A: Math. Gen. **36**, 3629 (2003).
- [43] T. Guhr and B. Kälber, J. Phys. A: Math. Gen. **36**, 3009 (2003).
- [44] R. Schäfer, N. F. Nilsson and T. Guhr, Quantitative Finance **10**, 107 (2010).
- [45] T. Schmitt, R. Schäfer, D. Wied and T. Guhr, Working paper (2013).
- [46] J. D. Noh, Phys. Rev. E **61**, 5981 (2000).
- [47] S. Pafka and I. Kondor, Physica A **319**, 487 (2003).
- [48] V. A. Marčenko and L. A. Pastur, Math. USSR Sb. **1**, 457 (1967).
- [49] A. Pandey and S. Ghosh, Phys. Rev. Lett. **87**, 024102 (2001); S. Ghosh and A. Pandey, Phys. Rev. E **65**, 046221 (2002).
- [50] J. Baik *et al.*, Ann. Probab. **33**, 1643 (2005); S. Péché, J. Multivariate Anal. **97**, 874 (2006).